



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MAINE

This label for SANDEA herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2020.

GROUP 2 HERBICIDE

Sandea® Herbicide

EPA Reg. No 81880-18-10163 EPA SLN NO. ME-160001B

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl, methyl 3-chloro-5-(4,6-dimethoxyimidin-2-ylcarbamoylsulfamoyl)	
-1-methylpyrazole-4-carboxylate.....	75.0%
OTHER INGREDIENTS	25.0%
	TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

PREHARVEST INTERVAL

The required days between last application and harvest are given in () after each crop name.

CROP	OZ/ACRE	DIRECTIONS FOR USE
13-07B LOWBUSH BLUEBERRIES (14)	1/2 - 1	<p>Apply uniformly with ground equipment in a minimum of 20 gal of water per acre. SANDEA should be tank mixed with products such as Velpar® Velossa (hexazinone ai's), or Sinbar® to broaden the spectrum of weeds controlled.</p> <ul style="list-style-type: none"> • Vegetative (Non-Crop) Year • Broadcast application prior to breaking dormancy in the Spring, or after blueberries are completely dormant in the Fall for control of labeled weeds. <p>Apply SANDEA as a single broadcast spray application. Applications applied 1 to 2 months prior to breaking dormancy will allow for better weed control.</p>
<p>PRECAUTIONS:</p> <ul style="list-style-type: none"> • Overlapping boom swaths increases the potential for phytotoxicity including leaf yellowing, reddening, and/or stunting • Consult "Use Precautions" and "For Optimum Results" of label for important usage information. • Preemergence applications of SANDEA when ground cover prevents contact with the soil will result in reduced or no residual activity. • SANDEA may not control ALS resistant weeds. <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not apply when frost is in the ground. • Do not apply to water saturated soils. • Do not apply to blueberries after vegetative bud break. • Do not apply to bushes established less than one year or to plants under stress. • Do not apply to areas where water is known to pond for periods of time following rainfall. • Do not apply SANDEA after the crop has progressed into budbreak or significant injury will occur. • Do not apply more than 1 application or 1 oz/A of product by weight (0.047 lb a.i./acre) per 12 month period. 		

24(c) Registrant: Gowan Company
P.O. Box 5569
Yuma, AZ 85366-5569